Pharmacy Bulletin

Pharmacy Department of Hospital Canselor Tuanku Muhriz

MONOFER (FERRIC DERISOMALTOSE AKA IRON ISOMALTOSIDE)

HCTM's NEW NON-FORMULARY & PARENTERAL IRON THERAPY DRUG

What is Monofer?

Active ingredient: Ferric derisomaltose Category: Non-Formulary

Price: Rm338.20/vial



Existing Parenteral Iron Therapy NEW!				
	Venofer	Cosmofer	Monofer	
Active ingredient	Iron Sucrose	Iron Dextrose	Ferric Derisomaltose	
mg/vial	100	100	500	
Price/vial	RM9.70	RM18.75	RM338.20	
Policy *For outpatient	For patients infused in HCTM ts, hospital cha	For Nephro patients, infused in HCTM rges apply	Must be bought from NF Pharmacy	

How is Monofer different from Venofer and Cosmofer?

Offers Convenience

- Single dose up to 20mg/kg no other dose limits
- No test dose required
- Infusion time of only 15 mins (for doses < 1g)

In Pregnancy & Lactation...

- Monofer should not be used during the first trimester. Iron deficiency anaemia occurring in the first trimester of pregnancy can, in most cases, be treated with oral iron.
- Monofer use should be confined to second and third trimesters.
- There is no available information on the excretion of Monofer in the human breast milk

Administration Comparison between Existing Parenteral Iron Therapy Drugs

	Venofer	Cosmofer	Monofer
Total dose infusion(TDI)	X	Ø	Ø
max single dose (per week)	7mg/kg or 500mg	20mg/kg	20mg/kg
Infusion time	TDI in 4-6hours	TDI in 4-6hours TDI in at least 15 mins (dose TDI in at least 30 mins (dose	
Test dose needed	Ø	Ø	*

Dose & Administration

The dose calculation is the same as for Venofer, Cosmofer & Monofer: Please refer to our formulary app

Monofer® should be added to 100ml sterile 0.9% sodium chloride. Do not dilute in more than 500ml.

EVERY PATIENT MUST BE OBSERVED FOR ADVERSE EFFECTS FOR AT LEAST 30 MINUTES

FOLLOWING EACH MONOFER INFUSION.

Google Play

App S

App S

The tables below provide information on the administration & preparation of Monofer® (iron isomaltoside 1000) for a range of doses.

Adia	Minimo	Monofer® (iron isomaltoside 1000)				
Minimum Iron dose administration			Maximum weekly dose = 20mg iron/kg body weight			
time	Dose	500mg	1,000mg	1,500mg	2,000mg	
≤1000mg	≤1000mg 20 minutes > 1000mg ≥30minutes	Dose (volume) of 100mg/ml vial	500mg (5ml)	1,000mg (10ml)	1,500mg (15ml)	2,000mg (20ml)
		Infusion fluid	100ml sodium chloride 0.9%*			
> 1000mg		Drug concentration	5mg/ml	10mg/ml	15mg/ml	20mg/ml

^{1.} Monofer® SPC – Electronic Medicines Compendium, November 2015.

^{2.} Sugimura M, Ohtani Y, Tamai S, et al. Ferric derisomaltose for the treatment of iron deficiency anemia with postpartum hemorrhage: Results of a single-arm, open-label, phase 3 study in Japan. J Obstet Gynaecol Res. 2023;49:946–55.

Prescribing ARBs separately



HCTM no longer purchases combination drug items of Angiotensin Receptor Blockers!
(except for Co-Diovan Tablet (Valsartan/Hydrochlorothiazide)

SO please indent ARBs separately!

Implication:

Once the drug is not available in the Formulary, the previous medication history will not list down the actual drug name but only a code will appear.

Action:

Specialist/ Doctors are requested to re-prescribe individual drug names so medication records can be recorded appropriately in the prescribing system.

500 1G TAB. (GLUCOPHAGE-G) : SELEPAS MAKAN A5V160 : SELEPAS MAKAN ara Pengambilan

A#

For example, if a patient was previously prescribed with EXFORGE 5mg/80mg tablets (Amlodipine 5mg/Valsartan 80mg), it now has to be indented separately: 1. Amlodipine 5 mg 1 tab OD 2. Valsartan 80mg 1 tab OD

> Drug class examples that were combined in one tablet



ARBs

 \rightarrow Irbesartan, Losartan, Telmisartan, Valsartan

CCBs

→ Amlodipine **Diuretics** → **Hydrochlorothiazide**

Please represcribe ARB combination drugs individually for accurate medication record:

- 1. CHECK if the patient is on any Combination **Drug containing ARB**
- 1. If a code (e.g T40A5) appears on the screen after copying and pasting the prescription, please **DELETE** the drug
- 1. PRESCRIBE the drugs again in the combination **SEPARATELY**
- (e.g Telmisartan 40mg/ Amlodipine $5mg \rightarrow$ Telmisartan 40mg on one line and Amlodipine 5mg on a different line).

Antihypertensive Combination Drugs Affected (with ARBs):

List of Combi Drugs containing ARBs	Drug combination content	Drugs that need to be prescribed separately	
Exforge	Amlodipine + Valsartan	AmlodipineValsartan	
Exforge HCT	Amlodipine+Valsartan+Hydrochl orothiazide (HCTZ)	AmlodipineValsartanHydrochlorothiazide	
Twynsta	Telmisartan + Amlodipine	TelmisartanAmlodipine	
Coaprovel	Irbesartan + HCTZ	IrbesartanHydrochlorothiazide	
Micardis Plus	Telmisartan + HCTZ	TelmisartanHydrochlorothiazide	
Hyzaar & Fortzaar	Losartan + HCTZ	LosartanHydrochlorothiazide	
Cozaar XQ	Amlodipine + Losartan	AmlodipineLosartan	

HCTM Formulary App is on...



